

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CHOICE TRIFLURALIN 480

Herbicide

ACTIVE CONSTITUENT: 480 g/L TRIFLURALIN
SOLVENT: 474 g/L LIQUID HYDROCARBON

GROUP D HERBICIDE

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as listed in the Directions for Use table.

APVMA APPROVAL No: 59371 / 0305

IMPORTANT: READ THIS LEAFLET BEFORE USE



GROW CHOICE PTY. LTD.

113 Fitzroy Street
Tamworth NSW 2340

Tel: 02 6766 3979

ACN 069 839 961

DIRECTIONS FOR USE

1. FIELD CROPS

Situation & Crop	Weeds	State	Rate L/ha Soil Type			Critical Comments
			Light	Medium	Heavy	
Chickpeas	Annual Ryegrass, Paradoxia Grass (Canary Grass), Wireweed (Hogweed), Black Pigweed, Suppression of Climbing Buckwheat (Black Bindweed), Wild Oats	Qld only	1.25 - 1.7L/ha	1.25 - 1.7L/ha	1.25 - 1.7L/ha	Use 1.25L/ha when applying immediately prior to sowing. Use 1.7L/ha when applying to dry soil before the planting rain.
		Vic only	800 mL/ha plus 1.6L/ha AvadexXtra			Incorporate as per recommendations for wheat, barley and triticale.
Adzuki Beans, Cowpeas, Lablab, Mung Beans, Borlotti Beans, Red Kidney Beans	Red & White Furnitory, Rough Poppy, Wireweed, Annual Ryegrass, Barley Grass, Canary Grass, Sand Fescue, Suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burrweed, Brome Grass, Cereal Oats and soil surrace Wild Oats	SA only	1.25L/ha	1.25L/ha	1.25L/ha	Apply to level seedbed 0 to 4 weeks before sowing. Incorporate as per Incorporation Table 6.
		NSW, ACT only	1.2 - 1.5L/ha	1.5L/ha	1.7L/ha	Apply from 4 weeks up to just prior to sowing. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.

Faba Beans	Annual Ryegrass, Barley Grass, Capeweed, Corn Gromwell (Sheepweed), Fumitories, Geranium, Ivy Leaf, Speedwell, Mustards, Turnips, Wireweed, Suppression of Bromo Grass, Soursob and Wild Oats	SA, WA only	800mL/ha plus 1.8-2L/ha Simazine 500SC	Apply to bare moist soil and grate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to ridged or excessively cloddy soil. For full reliable results, significant rainfall (20 or 30mm) is necessary within 2-3 weeks of application.
Pigeon Peas	Amaranthus, Barmyard Grass, Canary Grass, Crowsfoot Grass, Pigweed, Spiny Burrgrass, Summer Grass, Wild Oats, Wireweed (hogweed), Suppression of Yellow Vine (Caltrop), From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass	NSW, ACT only	1.2L/ha 1.5L/ha 1.7L/ha	Apply between 4 weeks and just before sowing. Refer to Incorporation Table 3, 4 or 6 for suitable method of incorporation.
Navy beans Soybeans	Annual Ryegrass, Barmyard Grass, Canary Grass, Caltrop (Bullhead, Yellow Vine), Crab Grass, Mossman River Grass (Innocent Grass) Pigweed, Redroot (Amaranthus) Redshank (Prince of Wales feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	All States	1.2L/ha 1.2L/ha 1.5L/ha 1.5L/ha 1.7L/ha 2.3L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Lentils	Annual Phalaris, Annual Ryegrass, Wild Oats, Wireweed Fumitory - Red and White, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue	NSW, ACT only SA only	800mL/ha 1.25L/ha 1.2L/ha 1.25L/ha	Apply 1 to 4 weeks before sowing. Apply 1 to 4 weeks before sowing.

Vetch	Annual Ryegrass, Deadnettle, Wireweed, soil surface Wild Oats, suppression of Brome Grass, Rough Poppy, Speedwell, Three Cornered Jack, Yellow Burr Weed, Sheepweed	SA, WA only	1.7L/ha	1.7L/ha	1.7L/ha	Apply to level seedbed 0 to 4 weeks before sowing. Refer to Incorporation Table 6 for method of incorporation.
Cotton	Annual Ryegrass, Barmyard Grass, Canary Grass, Catnip (Bulthead / Yellow Vine), Crab grass, Furnitory, Mossman Player Grass (Innocent Weed), Pigweed, Phalaris spp, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, soil surface Wild Oats, Winter Grass, Wireweed (Hogweed), Black Pigweed (Old only), From seed only, Columbus - Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	Old, NSW, ACT, WA only NSW, ACT, SA, WA, Vic, Tas only All States	1.2L/ha	1.7L/ha	2.3L/ha	Spray between 6 weeks and just before sowing takes place. Refer Incorporation Table 1 and 2 for method of incorporation. Autumn Sowing - Apply from 4 weeks to 7 days before sowing takes place. Refer Incorporation Table 6 for method of incorporation. Spring Sowing - Apply between 4 weeks and 3 days before sowing takes place. Refer Incorporation Table 6 for method of incorporation. In both cases seedling disease, cold weather, excessive moisture, high salt concentrations and drought could weaken crop seedlings and damage could occur from the use of this product. Temporary drop suppression could result.
Legume Seed Crop Establishment - Annual Medics - Clover Berseem, Red, Strawberry (Sub & White) Lucerne - for hay and seed crop			1.2L/ha	1.2L/ha	1.7L/ha	
Linseed		NSW, ACT, SA, WA, Vic only	1.2L/ha	1.5L/ha	1.7L/ha	Spray 2-4 weeks before sowing. Sowing depth should be 1.3 to 2.5cm. Deeper sowing may result in some stand reduction. Refer Incorporation Table 6 for method of incorporation.
Peanuts		WA, Old only				Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Peas		All States				Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 or 11 for method of incorporation.
Canola, Safflower						

Sugarcane Early season	Weeds as listed above	Qld, NSW, only	3.0L/ha	3.0L/ha	3.0L/ha	Apply to plant cane after emergence to "out of hand" stage. Apply to ratoon cane immediately after harvest. Refer Incorporation Table 3, 7 or 10 for suitable method of incorporation.
			2.3L/ha	2.3L/ha	2.3L/ha	
Late season	Weeds as listed above	All States	1.2L/ha	1.5L/ha	1.7L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for method of incorporation.
Sunflowers			Lupins	NSW, ACT, Vic, Qld only	800mL/ha plus 3L/ha of Simazine 500SC Herbicide	1.25L/ha plus 1-1.5L/ha of Simazine 500SC Herbicide
Sunflowers	Annual Grasses and Broadleaf Weeds	NSW, ACT, Vic, Qld only				
			As above plus suppression of Brome Grass	WA only	Rate for Yellow Sands. Refer to incorporation tables 11, 12 or 13.	
Sunflowers	As above plus suppression of Brome Grass	As above plus suppression of Brome Grass	1.25L/ha plus 2L/ha of Simazine 500SC Herbicide	1.25L/ha plus 2L/ha of Simazine 500SC Herbicide	Rate for all other soil types. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Application should not be made to rigid or excessively cloddy soil. For Simazine to be effective sufficient rainfall (20 to 30mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Results with Simazine can be variable if seasonal conditions are dry prior to sowing and Lupins are sown into dry or low moisture seed beds.	

Lupins	Capeweed, Doublegee, Wild Radish, Wild Turnip plus suppression of Annual Ryegrass, Wild Oats and Brome Grass	WA only	1.25L/ha plus 2L/ha of Diuron 500 Herbicide	DO NOT use on white or grey sands as severe crop damage may result. Use tank mix of Diuron and Trifluralin where Annual Ryegrass is present. Apply at pre-sowing stage when using incorporation method in Table 13. For post-sowing Pre-emergent application, ensure seed is adequately covered with soil. Refer to Incorporation Table 12.
	Red & White Fumitory, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue, suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr Weed, Brome Grass, Cereal Oats, soil surface Wild Oats	SA only	1.25 - 1.7L/ha 1.25 - 1.7L/ha 1.25 - 1.7L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 6 for method of incorporation.
Tobacco	Above weeds plus Capeweed, Common Fumitory, Geranium, Indian Hedge Mustard, Sheepweed, Shepherd Pursue, Toadrush, Turnips, suppression of Ice Plant and Soursob		1.25L/ha to 1.7L/ha plus 2 to 4L/ha of Simazine 500 SC Herbicide	Use a low volume boom applying 50- 100 litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5 cm just prior to sowing the crop. Incorporate within 4 hours of application. Do not apply to ridged soil.
	Summer Grass, Crowsfoot Grass, Red Natal Grass, Love Grass, Button Grass, Rhodes Grass, Pigweed	Qld only	800mL/ha 1.2L/ha	Apply to soil 3-4 weeks prior to transplanting. The longer period to be used for applications made during June and July. Incorporate to a depth of 10cm.
	Crowsfoot Grass	NSW, ACT only	800mL/ha 1.2L/ha 1.2L/ha	Apply to light sandy soil 14 to 21 days before transplanting. Do not incorporate to a depth greater than 6cm. Apply to loam (medium soil) 14 to 21 days before transplanting. DO NOT incorporate to a depth of greater than 6cm.

Wheat, Barley & Triticale 1. Pre-Sowing Only	Annual Ryegrass, Wireweed (Hogweed), Phalaris spp.	NSW, ACT, WA & Vic only	800mL/ha	800mL/ha	800mL/ha	Apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Incorporation Table 6 for method of incorporation.
	Fumitory Canary Grass As above except for Fumitory	WA only Vic only Vic only				
2. Pre-Sowing & Post-sowing (self mulching soils)						Pre-Sowing - Apply more than 4 weeks before sowing to prevent crop damage. Post Sowing - Apply within 2 days after sowing to well prepared seedbed. Refer Incorporation Table 6 for suitable method of incorporation.
Wheat	Annual Ryegrass, Paradoxa Grass (Canary Grass), soil surface Wild Oats, Wireweed (Hogweed)	Qld only	800mL/ha	800mL/ha	800mL/ha	On non self-mulching soils apply 1-4 weeks before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. On self mulching soils as above except apply more than 4 weeks before sowing to prevent crop damage. Refer Incorporation Table 6 for method of incorporation.
Barley						Apply to self mulching and non self mulching soils from 1-4 weeks before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. Refer Incorporation Table 6 for suitable method of incorporation.
Wheat, Triticale, Rye	Annual Ryegrass, Red & White Fumitory, Phalaris spp. Wireweed, suppression of Deadnetties, Rough Poppy, Yellow Burr Weed	SA only				Apply 1-4 weeks before sowing. Sowing depth should be at least 5 cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Incorporation Table 6 for method of incorporation.
Barley	As above for SA plus Sand Bronte Grass		1.25L/ha	1.25L/ha	1.25L/ha	
Wheat & Triticale only	Annual Phalaris	NSW, ACT only				800mL/ha plus 20g/ha Chlorsulfuron 750 WG Herbicide
						If possible spray and incorporate into the soil in one place within 4 hours of spraying. Delay may cause inferior weed control.

2. FIELD CROPS: FOR USE IN NO-TILL / MIN TILL CROPPING SYSTEMS, PRE-SOWING OR INCORPORATED BY SOWING (IBS) (Using Incorporation Table 13)

Situation & Crop	Weeds	State	Rate L/ha Soil Type			Critical Comments
			Light	Medium	Heavy	
Wheat, barley and triticale	Annual Ryegrass, Wireweed, Phalaris spp., Fumitory	WA, SA, V/c, NSW, ACT only	1.5L - 2L/ha			Use the higher rate on lighter sandy/loam soils. DO NOT use on heavy soils. Use with Knife/Blade point sowing equipment. Use the higher rate for heavier stubble coverage. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels. Refer Table 13 for method of incorporation. WA only; Application can occur 0- 24 hours prior to incorporation by sowing. For best results apply as close as possible to sowing (within 12 hours). Application 12-24 hours before sowing may be more adversely affected by above average soil moisture, warm temperatures and high weed seed densities. These factors individually or combined may reduce final weed control levels.
Chickpeas		WA only	1.25 - 1.7L/ha plus 2L/ha Simazine 500SC Herbicide			Incorporate as per incorporation Table 13.

2. VEGETABLES, ORCHARDS AND VINEYARDS

Situation & Crop	Weeds	State	Rate L/ha Soil Type			Critical Comments
			Light	Medium	Heavy	
Transplants only Broccoli, Cabbage, Cauliflowers, Tomatoes	Annual Ryegrass, Barnyard Grass, Canary Grass, Calltrop, (Bullhead Yellow Vine), Crab Grass, Mossman River Grass (Innocent Grass) Pigweed Redroot (Amaranthus) Redshank (Prince of Wales feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed)	All States	1.2L/ha	1.7L/ha	2.3L/ha	Spray between 4 weeks and just before sowing takes place. Refer Incorporation Table 3, 4 or 5 for suitable method of incorporation.
Direct Seeded Only Broccoli Brussels Sprouts, Cabbage		Vic, Qld only				
Cauliflower		All States				
Carrots	From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	Vic only				
Chicory		All States	1.2L/ha	1.5L/ha	1.7L/ha	
Green Beans						
Orchards and Vineyards		Qld, SA, WA, Vic, Tas only	1.2L/ha	1.7L/ha	2.3L/ha	Apply to new planting during pre plant cultivation. Apply to established crops in spring after weeds and green manure crop has been ploughed into ground. Refer Incorporation Table 8 or 9 for suitable method of incorporation.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

THIS PRODUCT MUST BE INCORPORATED INTO THE SOIL WITHIN 4 HOURS OF APPLICATION EXCEPT IN WESTERN AUSTRALIA WHERE THE CROPS SOWN WITH MINIMUM TILLAGE SOWING EQUIPMENT (FITTED WITH KNIFE POINTS OR BLADES LESS THAN 12MM WIDE, USUALLY WITH PRESS WHEELS) WHERE APPLICATION MAY OCCUR UP TO 24 HOURS BEFORE INCORPORATION BY THE SOWING PROCESS.

INCORPORATION TABLE

1. Prior to furrowing out: 2 workings at an angle required using Offset or Tandem disc harrows.
2. After furrowing out: 2 workings required using Go-Devil discs or Lilliston cultivators set at 10cm depth.
3. Rotary Hoe: 1 working required at 5-7.5cm depth.
Sugar Cane: 7.5-13cm depth.
4. Offset or Tandem Disc Harrows: (preferably with spiked harrows in tandem) 2 workings at an angle required at 7.5-15cm depth at 6.5-10km per hour.
5. Heavy Diamond or Stump Jump Harrows - (weighted 20-30kg per section) at 10-13km/h speed. Then cross work with offset or tandem disc harrows set to 7.5-15cm depth at speed 6.5-10km per hour.
6. Weighted Heavy Diamond or Stump Jump Harrows (weighted with 20-30kg per section) at 10-13km per hour. Cross work with combine at 5-7.5cm depth at speed of 10-13km/hr.
7. Disc Ratoon Cultivator: 2 workings needed with discs and cultivator set at 7.5-13cm depth.
8. Offset or Tandem Disc Harrows: set at 7.5-15cm depth. A second discing is required working in opposite direction with discs set to throw treated soil into tree or vine row.
9. Rotary Hoe: 1 working needed at 5-10cm depth.
10. Offset Discs (Bumpers): 2 workings needed at depth of 7.5-13cm.
11. Incorporation by Sowing (IBS) on suitably prepared seedbed with heavy diamond harrows trailing or as a separate operation.

12. Post-sowing pre-emergence: Use heavy diamond harrows cross working at right angles to the direction of sowing. Do not attempt this method of incorporation on poorly prepared, clumpy or cloddy soils.
13. Incorporation by sowing (IBS) with knife or blade points. Use press wheels to avoid dragging treated soil back into the seed furrow. Maintain slow to moderate speed to ensure that soil throw is not into adjacent furrows. Note: Knife or blade point systems can result in poor weed control in the seed furrow as chemical displacement from this zone occurs. Stubble coverage above 40-50% ground cover can reduce weed control below acceptable levels.

A Knife or blade point is 12mm or less, has no inverted wings, inverted T or blade, and is generally placed on a minimum 8 inch tyne spacing.

MIXING

This product is an emulsifiable concentrate which mixes readily with water. Add the recommended amount to the spray tank during filing operation and apply 70-450L of water/ha (broadcast basis) dependent on soil type and stubble coverage level. For minimum tillage/stubble retention seeding systems use of the higher water volumes may help reduce the impact of stubble.

Ensure adequate agitation is continued throughout the operation. Leaving the made up spray mixture for long periods of time without agitation is not recommended.

Under hot conditions or where possible spray and incorporate into the soil in one operation. Delay may cause inferior weed control. Use properly calibrated standard low pressure (170-340 kilopascal) boom type sprayer with fan tips.

CONDITIONS FOR BEST RESULTS

This product must be thoroughly incorporated as recommended. Soil should be well worked and free of weeds at time of application. Product effectiveness may be reduced by inadequate incorporation, high organic matter, excess clods, crop or trash residues, stones or other foreign matter and in areas of unnaturally high weed seed population such as header tracks or livestock rest areas. Trifluralin is volatile and disappears from exposed surfaces. Loss is hastened by high temperatures, winds or warm moist soil.

INTEGRATED WEED MANAGEMENT

The use of Integrated Weed Management techniques in conjunction with Choice Trifluralin are always recommended. Agronomic practices that reduce the weed seed bank in the soil prior to the use of Choice Trifluralin will result in higher weed control levels from Choice Trifluralin. Failure to use Agronomic and Integrated Weed Management practices that reduce the weed seed bank in the soil will result in higher weed seed populations. Paddocks with excessively high weed seed banks may have sufficient weed numbers surviving such that final weed control may be considered below a commercially acceptable level and additional herbicide treatments may be necessary. The use of Integrated Weed Management techniques will also reduce potential for the development or survival of Group D herbicide resistant weed biotypes.

WILD OATS

Germinating wild oat seeds lying on soil surface will not be controlled. Therefore, specific wild oat control is only possible with shallow cultivation. Poor control will occur on self mulching soils and all soil types where deep cultivation is practiced.

COMPATIBILITY

This product may be mixed in the spray tank with: Herbicides: Avadex Xtra, Chlorsulfuron, Triasulfuron, Diuron Granules (900g/kg diuron), Diuron 500 Flowable, Simazine 500 Flowable, Simazine 900 WG, Nuquat, Gramoxone, Sprayseed, Broadstrike, Spinnaker, Bladex, Metribuzin (750g/kg), Insecticides: Chlorpyrifos 500 EC, Endosulfan 350EC.

- NOTE:
1. Information on compatibility is understood to be correct at the time of publication, however products may vary from time to time, therefore a small scale compatibility test should be carried out before mixing in the spray tank.
 2. Observe any mixing sequence instructions for tank mix products.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible, otherwise, spray tanks, pumps, lines and nozzles should be thoroughly rinsed several times with clean water following application. Spraymate Tank & Equipment Cleaner is suitable for this purpose and will also remove Trifluralin stains.

RESISTANT WEEDS WARNING

GROUP **D** HERBICIDE

Choice Trifluralin 480 Herbicide is a member of the Dinitronilines group of herbicides. Choice Trifluralin 480 Herbicide has the inhibitors or tubulin formation mode of action. For weed resistance management Choice Trifluralin 480 Herbicide is a Group D herbicide.

Some naturally occurring weed biotypes resistant to Choice Trifluralin 480 Herbicide and other group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Choice Trifluralin 480 Herbicide or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use Grow Choice Pty Ltd accepts no liability for any losses that may result from the failure of Choice Trifluralin 480 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT use in high winds.

DO NOT exceed rates specified, to avoid crop damage.

DO NOT plant sensitive grasses such as oats, sorghum, millets, phalaris spp, ryegrass or wheat for 12 months following the use of this product except where wheat follows wheat or other winter crops.

DO NOT plant oilseed poppies when a detectable residue of Trifluralin is present in the soil. Levels as low as 0.02 ppm may interact with other unfavourable factors (moisture, stress disease etc) to reduce poppy growth and vigour.

DO NOT apply to orchards and vineyards after first flush of growth or when residues can lodge on or in fruit. Reduced germination of wheat and barley may occur due to combination of following circumstances and use of this product:

- Short Coleoptile cultivars.
- Use of seed dressings (except Vitavax).
- Shallow or uneven seedling depth.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants adjacent crops, crop lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area away from children, animals, food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers (1000L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the prepared spray wear cotton overalls, buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

For further information refer to Material Safety Data Sheet which can be obtained from the supplier.

NOTICE TO BUYER

Grow Choice Pty. Ltd. will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.

