

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PROMPT 750 WG

HERBICIDE

ACTIVE CONSTITUENT:

750 g/kg TRIBENURON-METHYL

GROUP B HERBICIDE

For the control of certain broadleaf weeds in fallows
and pre-cropsituations as per the
DIRECTIONS FOR USE table.

**IMPORTANT: Read the attached leaflet before
using this product**

APVMA Approval No: 81740/103933



GROW CHOICE PTY LTD

113 Fitzroy Street
Tamworth NSW 2340

Tel: (02) 6766 3979

ACN 161 264 884

DIRECTIONS FOR USE:

RESTRAINTS

DO NOT apply to weeds that are not actively growing or stressed by any cause such as adverse weather conditions, drought and water logging. Broadleaf weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or only growth suppression. If heavy grazing has occurred allow regrowth before spraying. **DO NOT** apply if rainfall is expected within 2 hours.

Minimum volumes of prepared spray to be applied:

- **Ground application** - 50L/ha
- **Aerial application** - 20L/ha

Use prepared spray suspensions within two days of preparation, otherwise significant breakdown may occur.

Prompt 750 WG Herbicide used alone – NSW & Qld Only

CROP	WEEDS CONTROLLED	RATE g/ha	WEEDS GROWTH STAGE AT APPLICATION	CRITICAL COMMENTS
FALLOW and PRE-CROP (- refer to Crop Options section for minimum intervals between application and sowing)	Amaranthus / Boggabri weed (<i>Amaranthus mitchelli</i>)	25	Apply up to the 10-leaf stage	As a herbicide Resistance countering measure, DO NOT use an ALS inhibitor herbicide in the crop following the use of PROMPT 750 WG Herbicide as a preceding fallow or pre-crop treatment.
	Caltrop / Yellowvine (<i>Tribulus terrestris</i>)			
	Sowthistle / Milk Thistle (<i>Sonchus oleraceus</i>)			
	Deadnettle (<i>Lamium amplexicaule</i>)	25	Apply up to the 6-leaf stage	As a herbicide Resistance countering measure, DO NOT use an ALS inhibitor herbicide in the crop following the use of PROMPT 750 WG Herbicide as a preceding fallow or pre-crop treatment. If weeds are at a more advanced growth stage and/or are present in high numbers (greater than 50 per m ²) use a Glyphosate mixture as specified in the PROMPT 750 WG Herbicide and Glyphosate Tank Mixes table .

CROP	WEEDS CONTROLLED	RATE g/ha	WEEDS GROWTH STAGE AT APPLICATION	CRITICAL COMMENTS
FALLOW and PRE-CROP (- refer to Crop Options section for minimum intervals between application and sowing)	Medics (Native) <i>(Medicago spp.)</i>	30	Apply up to 5cm diameter	If weeds are at a more advanced growth stage and/or are present in high numbers (greater than 50 per m ²) use a Glyphosate mixture as specified in the PROMPT 750 WG Herbicide and Glyphosate Tank Mixes table .
	New Zealand Spinach <i>(Tetragonia tetragonoides)</i>	20	Apply up to the 10-leaf stage	
	Prickly Lettuce <i>(Lactuca serriola)</i>	30	Apply up to the 4-leaf stage	If weeds are at a more advanced growth stage and/or are present in high numbers (greater than 50 per m ²) use a Glyphosate mixture as specified in Tank Mixes table .
Always add non-ionic surfactant (1000g/L – non buffering type) at 100 mL/100L (0.1% v/v) of final volume. When applying PROMPT 750 WG Herbicide alone by air, in addition to a non-ionic surfactant, add a mineral spray oil at 1L/100L (1% v/v) of final spray volume.	Turnip Weed <i>(Raphistrum rugosum)</i>	20	Apply up to flowering	

Prompt 750 WG Herbicide + Glyphosate Tank Mixes Table – NSW & Qld only
 For the control of these weeds in addition to those in the Prompt 750 WG Herbicide alone table

CROP	WEEDS CONTROLLED	RATE g/ha	WEEDS GROWTH STAGE AT APPLICATION	CRITICAL COMMENTS
FALLow and PRE-CROP (- refer to Crop Options section for minimum intervals between application and sowing)	Black Bindweed (<i>Fallopia convolvulus</i>)	25 plus Glyphosate at Rate 1*	Apply up to the 10-leaf stage	For best control, apply to small actively growing weeds. Larger weeds are more difficult to control.
	Common Thornapple (<i>Datura stramonium</i>)	20 plus Glyphosate at Rate 2*	Apply up to the 8-leaf stage	
	Deadnettle (<i>Lamium amplexicaule</i>)	25 plus Glyphosate at Rate 2*	Apply up to the 10-leaf stage	
	Mintweed (<i>Salvia reflexa</i>)	15 plus Glyphosate at Rate 1*	Apply up to the 6-leaf stage	
	Pigweed / Portulaca (<i>Portulaca oleracea</i>)	20 plus Glyphosate at Rate 1*		
	Prickly Lettuce (<i>Lactuca serriola</i>)			

*Glyphosate Rates

Rate 1 : Choice Glyphosate 700 Soluble Granular Herbicide @ 385g/ha or Glyphosate 450g/L @ 600mL/ha

Rate 2 : Choice Glyphosate 700 Soluble Granular Herbicide @ 260g/ha or Glyphosate 450g/L @ 400mL/ha

Always add non-ionic surfactant (1000g/L – non buffering type) at 100 mL/100L (0.1% v/v) of final volume.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

Best results are obtained when **PROMPT 750 WG Herbicide** is applied to young actively growing weeds. The rate will depend on weed spectrum at time of application. The degree of control and duration of effect are dependent on rate used, sensitivity and size of target weeds and environmental conditions at the time of and following application.

PROMPT 750 WG Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms of dying weeds (discolouration) may not be noticeable for 1 to 3 weeks after application depending on the environmental conditions and susceptibility. Warm, moist conditions following treatment promotes the activity of Choice **PROMPT 750 WG Herbicide**, while cold, dry conditions delay activity. Weeds hardened-off by cold weather and/or drought stress will be less susceptible. Where recommended the addition of Glyphosate will enhance knockdown and control of weeds. It also helps minimise the risk of resistant weeds occurring.

CROP SAFETY

DO NOT apply chlorsulfuron, metsulfuron-methyl or triasulfuron products in a tank mix with **PROMPT 750 WG Herbicide** as a fallow or pre-crop treatment.

SPRAY PREPARATION

PROMPT 750 WG Herbicide is a dry flowable formulation to be mixed with water and applied as a spray. Partially fill the spray tank with water. Measure the amount of **PROMPT 750 WG Herbicide** required for the area to be sprayed.

Add the correct amount of **PROMPT 750 WG Herbicide** to the spray tank with the agitation system engaged. Top up to the correct volume with water. The material must be kept in suspension at all times by continuous agitation. In tank mixes, **PROMPT 750 WG Herbicide** must be in suspension before adding the companion herbicide or surfactant/wetting agent.

COMPATIBILITY

PROMPT 750 WG Herbicide is compatible with Glyphosate herbicides.

USE OF SURFACTANT/WETTING AGENT

Always add non-ionic surfactant (1000g/L – non-buffering type) at 100 mL/100L (0.1% v/v) of final spray volume. When applying **PROMPT 750 WG Herbicide** alone by air in addition to a non-ionic surfactant add a mineral spray oil at 1L/100L (1% v/v) of final spray volume.

GROUND SPRAYING

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping. Apply a minimum of 50L prepared spray/ha.

AERIAL SPRAYING

Apply a minimum of 20L/ha water. Application in 30L/ha will improve performance reliability. Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops or fallow areas likely to be planted to these crops - see Crop Options section. Turn off spray boom whilst passing over creeks and dams.

DO NOT apply when a temperature air inversion is likely to occur.

SPRAYER CLEANUP

To avoid subsequent injury to crops immediately after spraying thoroughly remove all traces of **PROMPT 750 WG Herbicide** from mixing and spray equipment as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through boom and hoses using at least one-tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water adding household ammonia to give a 1% solution (equivalent ratios include 1/3 litre household ammonia product, containing 9.5% ammonia) for each 100 litres of tank volume. Agitate, and then flush the boom and hoses with the cleansing solution.

Top up with water making sure the tank is completely full and allow standing for 15 minutes with agitation engaged. Again flush the boom and hoses and drain tank completely.

4. Nozzles and screens should be removed and cleaned separately with ammonia solution at the same concentration as used for the sprayer.

5. Rinse the tank with clean water and flush through boom and hoses using at least one-tenth of the spray tank volume. Drain tank completely.

6. **DO NOT** spray the tank rinsate onto sensitive crop or land intended for cropping with sensitive crop.

NOTE: IF not possible to drain the tank completely, step 3 must be repeated before going on to step 4.

CROP OPTIONS

Land previously treated with **PROMPT 750 WG Herbicide** may be sown to any of the specified crops after the interval indicated in the following table:

Minimum Interval	Crops (Note 2)
3 days	Barley, Oats, Wheat
7 or 21 days (Note 1)	Maize, Mungbeans, Sorghum, Soybeans, Sunflowers

Note 1:

PROMPT 750 WG Herbicide is broken down in soil, primarily by chemical hydrolysis, but to a lesser degree by microbial degradation. Breakdown is fastest in warm, wet acid soils and slower in cold, dry alkaline soils. For the above summer crops, if minimum soil temperatures at planting depth are less than or equal to 15°C for three consecutive days, then plant back intervals should be extended to 21 days.

Note 2:

As a herbicide resistance countering measure, **DO NOT** use a Group B Herbicide in any of these crops following the use of **PROMPT 750 WG Herbicide** alone as a preceding fallow or pre-crop treatment. Tolerance of other crops (growth through to maturity) should be determined on a small scale before re-sowing into larger areas.

RESISTANT WEEDS WARNING

GROUP	B	HERBICIDE
--------------	----------	------------------

PROMPT 750 WG Herbicide is a member of the Sulfonylurea group of herbicides. **PROMPT 750 WG Herbicide** has the Inhibitor of Acetolactate Synthase (ALS) mode of action. For weed resistance management **PROMPT 750 WG Herbicide** is a Group B Herbicide. Some naturally occurring weed biotypes resistant to **PROMPT 750 WG Herbicide**

and other Group B Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by **PROMPT 750 WG Herbicide** or other Group B Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Grow Choice Pty Ltd accepts no liability for any losses that may result from failure of **PROMPT 750 WG Herbicide** to control resistant weeds. Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors. DO NOT use Group B Herbicide against the same weed in the crop following the use of **PROMPT 750 WG Herbicide** alone either as a fallow or pre-crop treatment. This does NOT apply if the fallow or pre-crop use if **PROMPT 750 WG Herbicide** was tank mixed with Glyphosate. Avoid the prolonged use of Group B Herbicide on the same weed population. If the user suspects that an ALS inhibitor resistant weed is present, **PROMPT 750 WG Herbicide** or other Group B Herbicides recommended for the control of those weeds should not be used. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Grow Choice Pty Ltd representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGETED PLANTS INCLUDING AQUATIC PLANTS.

Injury to or loss of desirable trees, vegetation including aquatic plants may result from failure to observe the following:

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

The use pattern is such that a withholding period for harvest or grazing is not required. It is recommended however, not to graze treated areas for 2 to 3 days to ensure adequate control is achieved.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. DO NOT re-use container.

Triple-rinse containers before disposal. Dispose of rinsate or any undiluted chemical according to state/territory legislative requirements.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When preparing spray wear elbowlength PVC gloves and face shield or goggles. If product in eyes, wash out immediately with water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. **Phone Australia 13 11 26.**

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier.

NOTICE TO BUYER

Grow Choice Pty. Ltd. will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.